

# IBSRELA<sup>®</sup> (tenapanor) tablets

## An Innovative Approach for the Treatment of Irritable Bowel Syndrome With Constipation (IBS-C) in Adults

We invite you to join us for this presentation that will review:

- IBS-C multifactorial pathophysiology
- IBSRELA information, including
  - First-in-class novel mechanism of action
  - Efficacy and safety based on two Phase 3 clinical trials

The Product Theater content and views expressed therein are those of the sponsor and not of Digestive Disease Week<sup>®</sup>



### Brooks D. Cash, MD, FACG, AGAF, FASGE

Dan and Lillie Sterling Professor of Medicine  
Chief, Gastroenterology, Hepatology, and Nutrition  
UT Health Science Center at Houston  
McGovern Medical School  
Houston, TX



### Mark Pimentel, MD, FRCP(C)

Executive Director  
Medically Associated Science and Technology (MAST) Program  
Cedars-Sinai Medical Center  
Los Angeles, CA

**MONDAY  
MAY 23**

**2:30 - 3:15 PM  
Hall F2**

## IMPORTANT SAFETY INFORMATION

### **WARNING: RISK OF SERIOUS DEHYDRATION IN PEDIATRIC PATIENTS**

IBSRELA is contraindicated in patients less than 6 years of age; in nonclinical studies in young juvenile rats administration of tenapanor caused deaths presumed to be due to dehydration. Avoid use of IBSRELA in patients 6 years to less than 12 years of age. The safety and effectiveness of IBSRELA have not been established in patients less than 18 years of age.

### **CONTRAINDICATIONS**

- IBSRELA is contraindicated in patients less than 6 years of age due to the risk of serious dehydration.
- IBSRELA is contraindicated in patients with known or suspected mechanical gastrointestinal obstruction.

### **WARNINGS AND PRECAUTIONS**

#### **Risk of Serious Dehydration in Pediatric Patients**

- IBSRELA is contraindicated in patients below 6 years of age. The safety and effectiveness of IBSRELA in patients less than 18 years of age have not been established. In young juvenile rats (less than 1 week old; approximate human age equivalent of less than 2 years of age), decreased body weight and deaths occurred, presumed to be due to dehydration, following oral administration of tenapanor. There are no data available in older juvenile rats (human equivalent 2 years to less than 12 years).
- Avoid the use of IBSRELA in patients 6 years to less than 12 years of age. Although there are no data in older juvenile rats, given the deaths in younger rats and the lack of clinical safety and efficacy data in pediatric patients, avoid the use of IBSRELA in patients 6 years to less than 12 years of age.

### **Diarrhea**

Diarrhea was the most common adverse reaction in two randomized, double-blind, placebo-controlled trials of IBS-C. Severe diarrhea was reported in 2.5% of IBSRELA-treated patients. If severe diarrhea occurs, suspend dosing and rehydrate patient.

### **MOST COMMON ADVERSE REACTIONS**

The most common adverse reactions in IBSRELA-treated patients (incidence  $\geq 2\%$  and greater than placebo) were: diarrhea (16% vs 4% placebo), abdominal distension (3% vs  $<1\%$ ), flatulence (3% vs 1%) and dizziness (2% vs  $<1\%$ ).

### **INDICATION**

IBSRELA (tenapanor) is indicated for the treatment of Irritable Bowel Syndrome with Constipation (IBS-C) in adults.

**Please see full Prescribing Information, including Boxed Warning, at [IBSRELA-hcp.com](http://IBSRELA-hcp.com).**

In accordance with the PhRMA Code on Interactions with Healthcare Professionals, this program is intended only for healthcare professionals who will find its content clinically relevant to their practice. Accordingly, non-healthcare professionals (ie, spouses and other guests) may not attend. Invitation is extended only to those HCPs whose attendance would not violate compliance with state law requirements in which such HCP is licensed to practice.

