

VOWST®

(fecal microbiota spores,
live-brpk) capsules

VOWST is the **FIRST** and **ONLY** orally administered
microbiome therapeutic to prevent *C. difficile** recurrence^{1,2}

SUPERIOR PREVENTION OF *C. DIFF* RECURRENCE VS ANTIBIOTICS ALONE

PRIMARY ENDPOINT

88%
P<0.001

RECURRENCE FREE AT
8 WEEKS

(vs 60% of participants
receiving antibiotics alone)^{1,2,†}

SECONDARY ENDPOINT

79%
P<0.001

RECURRENCE FREE AT
24 WEEKS

(vs 53% of participants
receiving antibiotics alone)^{1,3,‡}

GENERALLY WELL TOLERATED THROUGH 8 WEEKS¹

- The majority of adverse reactions were mild or moderate in severity
- Most adverse reactions occurred within 10 days of starting VOWST
 - The median duration of these events was ≤5 days
- No serious adverse events were considered related to the use of VOWST

IMPORTANT SAFETY INFORMATION

WARNINGS AND PRECAUTIONS

Transmissible infectious agents: Because VOWST is manufactured from human fecal matter, it may carry a risk of transmitting infectious agents. Report any infection that is suspected to have been transmitted by VOWST to Aimmune Therapeutics, Inc. at 1-833-246-2566.

Potential presence of food allergens: VOWST may contain food allergens. The potential to cause adverse reactions due to food allergens is unknown.

ADVERSE REACTIONS

The most common adverse reactions (reported in ≥5% of VOWST-treated participants, and at a rate greater than placebo) were abdominal distension (31.1%), fatigue (22.2%), constipation (14.4%), chills (11.1%), and diarrhea (10.0%).

To report SUSPECTED ADVERSE REACTIONS, contact Aimmune Therapeutics at 1-833-AIM-2KNO (1-833-246-2566), or the FDA at 1-800-FDA-1088, or visit www.fda.gov/MedWatch.

DRUG INTERACTIONS

Do not administer antibacterials concurrently with VOWST.

INDICATION

VOWST is indicated to prevent the recurrence of *Clostridioides difficile* infection (CDI) in individuals 18 years of age and older following antibacterial treatment for recurrent CDI (rCDI).

Limitation of Use: VOWST is not indicated for treatment of CDI.

Please see full Prescribing Information and Patient Information at booth #3041.

VISIT VOWST AT BOOTH #3041



Scan the QR or visit VOWSThcp.com to learn more

**Clostridioides difficile* (*C. diff*) infection.

†Calculated by the rate of recurrence, which was significantly lower in VOWST recipients compared to those receiving antibiotics alone—12% vs. 40%, respectively—a 0.32 relative risk (95% confidence interval [CI], 0.18, 0.58; P<0.001).^{1,2}

‡Relative risk of 0.46 (95% CI, 0.30, 0.73).^{1,3}

ECOSPOR III was a Phase 3, multicenter, double-blind, randomized, placebo-controlled trial evaluating *C. diff* recurrence at 8 weeks; secondary endpoints at 4, 12, 24 weeks. In the intent-to-treat population, participants received VOWST or placebo after standard-of-care antibiotic treatment (vancomycin/ fidaxomicin) and laxative. Those lost to follow-up, terminated prematurely, or died prior to end of study time interval counted as a recurrence. Data were rounded to the nearest whole number for presentation.^{1,3}

REFERENCES: 1. VOWST. Prescribing information. Seres Therapeutics, Inc.; 2023. Accessed February 9, 2024. https://www.serestherapeutics.com/our-products/VOWST_PI.pdf 2. Feuerstadt P, Louie TJ, Lashner B, et al. SER-109, an oral microbiome therapy for recurrent *Clostridioides difficile* infection. *N Engl J Med*. 2022;386:220-229. doi:10.1056/NEJMoa2106516 3. Cohen SH, Louie TJ, Sims M, et al. Extended follow-up of microbiome therapeutic SER-109 through 24 weeks for recurrent *Clostridioides difficile* infection in a randomized clinical trial. *JAMA*. 2022;328(20):2062-2064. doi:10.1001/jama.2022.16476

SERES and the Seres logo, VOWST and the VOWST logo are trademarks of Seres Therapeutics, Inc. Unless otherwise indicated, all trademarks are owned by Société des Produits Nestlé S.A., Vevey, Switzerland or used with permission.

© 2024 Nestlé. VOW-PM-USA-0332 03/24

